arXiv:1301.0699v2 [math.CA] 22 May 2013

CONVEXITY PROPERTIES OF GENERALIZED TRIGONOMETRIC AND HYPERBOLIC FUNCTIONS

ÁRPÁD BARICZ, BARKAT ALI BHAYO, AND RIKU KLÉN

Abstract. We study the power mean inequality for the generalized trigonometric and hyperbolic functions with two parameters. The generalized p-trigonometric and (p,q)-trigonometric functions were introduced by Lindqvist and Takeuchi, respectively.

1. **Introduction and Main Results**

The generalized trigonometric functions were introduced by Lindqvist [17] two decades ago. These p-trigonometric functions, $p > 1$, coincide with the usual trigonometric functions for $p = 2$. Recently, the p-trigonometric functions have been studied extensively, see for example [5, 8, 11, 16] and their references. Drábek and Manásevich [11] considered a certain (p,q)-eigenvalue problem with the Dirichlét boundary condition and found the complete solution to the problem. The solution of a special case is the function $\sin_{p,q}$, which is the first example of so called (p,q)-trigonometric function. Motivated by the (p,q)-eigenvalue problem Takeuchi [19] introduced the (p,q)-trigonometric functions, $p, q > 1$. These (p,q)-trigonometric functions have recently been studied also in [6, 7, 9, 10, 12], and the functions agree with the p-trigonometric functions for $p = q$.

The *Gaussian hypergeometric function* is the analytic continuation to the slit plane $\mathbb{C} \setminus [1,\infty)$ of the series

$$F(a,b;c;z) = {}_2F_1(a,b;c;z) = \sum_{n\geq 0} \frac{(a,n)(b,n)}{(c,n)} \frac{z^n}{n!}, \qquad |z| < 1,$$

for given complex numbers a, b and c with $c \neq 0, -1, -2, \ldots$. Here $(a,0) = 1$ for $a \neq 0$, and (a,n) is the *shifted factorial function* or the *Appell symbol*

$$(a,n) = a(a+1)(a+2)\cdots(a+n-1)$$

for $n = 1, 2, \ldots$. The hypergeometric function is used to define the (p,q)-trigonometric functions and it has numerous special functions as its special or limiting cases [1].

For $p, q > 1$ and $x \in (0,1)$ the function $\mathrm{arcsin}_{p,q}$ is defined by

$$\mathrm{arcsin}_{p,q}(x) = \int_0^x (1-t^q)^{-1/p} dt = x\, F\left(\frac{1}{p}, \frac{1}{q}; 1 + \frac{1}{q}; x^q\right).$$

We also define for $x \in (0,1)$ $\mathrm{arccos}_{p,q}(x) = \mathrm{arcsin}_{p,q}((1-x^p)^{1/q})$ (see [12, Prop. 3.1]) and for $x \in (0,\infty)$

$$\begin{aligned} \mathrm{arsinh}_{p,q}(x) &= \int_0^x (1+t^q)^{-1/p} dt = x\, F\left(\frac{1}{p}, \frac{1}{q}; 1 + \frac{1}{q}; -x^q\right) \\ &= \left(\frac{x^p}{1+x^q}\right)^{1/p} F\left(1, \frac{1}{p}, 1 + \frac{1}{q}; \frac{x^q}{1+x^q}\right). \end{aligned}$$

Their inverse functions are given by

$$\sin_{p,q}, \cos_{p,q} : (0, \pi_{p,q}/2) \to (0,1), \quad \sinh_{p,q} : (0,\infty) \to (0,\infty),$$

`File: PM_pqfuncs20130522.tex, printed: 2013-012-12, 3.46`

2010 *Mathematics Subject Classification.* 33C99, 33B99.

Key words and phrases. Power mean, eigenfunctions of p-Laplacian, generalized trigonometric functions, generalized inverse trigonometric functions, convexity with respect to power means, geometrical convexity.

where $\pi_{p,q}/2 = \operatorname{arcsin}_{p,q}(1)$. Now, we define $\tan_{p,q} : (0, \pi_{p,q}/2) \to (0, \infty)$ by

$$\tan_{p,q}(x) = \frac{\sin_{p,q}(x)}{\cos_{p,q}(x)}, \quad \cos_{p,q}(x) \neq 0,$$

and we denote its inverse by $\arctan_{p,q}$.

We introduce next a geometric definition for the functions $\sin_{p,q}$ and $\cos_{p,q}$. The definition is based on the formula [12, eq. (2.7)]

$$|\sin_{p,q}(x)|^q + |\cos_{p,q}(x)|^p = 1. \tag{1.1}$$

The usual trigonometric functions are geometrically defined by the unit circle. In the same manner we can define the (p,q)-trigonometric functions. Instead of the unit circle we use the Lamé-curve (or generalized superellipse) defined by

$$C = \{(x,y) \in \mathbb{R}^2 \colon x = \cos^{2/p} t,\ y = \sin^{2/q} t,\ t \in [0, \pi/2]\}.$$

Because $(\sin^{2/q} t)^q + (\cos^{2/p} t)^p = 1$ it is clear by (1.1) that $C = D$ for

$$D = \{(x,y) \in \mathbb{R}^2 \colon x = \sin_{p,q}(t),\ y = \cos_{p,q}(t),\ t \in [0, \pi_{p,q}/2]\}.$$

The curve C is defined only in the first quadrant, but by reflections we can easily extend C to form a circular curve, see Figure 1.



FIGURE 1. Left: Curve C for $(p,q) = (4,3)$ and points $(0, \sin_{4,3} x)$, $(\cos_{4,3} x, 0)$ and $(\cos_{4,3} x, \sin_{4,3} x)$. Right: Curve C for $(p,q) = (4,3)$ extended to all four quadrants (black) and the unit circle (gray).

Now, we recall the definition of convex functions with respect to power means. For $a \in \mathbb{R}$ and $x, y > 0$, the power mean M_a of order a is defined by

$$M_a(x,y) = \begin{cases} \left(\dfrac{x^a + y^a}{2}\right)^{1/a}, & a \neq 0, \\ \sqrt{x\,y}\,, & a = 0\,. \end{cases}$$

We consider the function $f : I \subset (0,\infty) \to (0,\infty)$, which is continuous, and let $M_a(x,y)$ and $M_b(x,y)$ be the power means of order a and b of $x > 0$ and $y > 0$. For $a, b \in \mathbb{R}$ we say that f is M_aM_b-convex (concave) or simply (a,b)-convex (concave), if for those a and b we have

$$f(M_a(x,y)) \leq (\geq) M_b(f(x), f(y)) \ \text{ for all } x, y \in I\,.$$

It is important to mention here that $(1,1)$-convexity means the usual convexity, $(1,0)$ means the log-convexity and $(0,0)$-convexity is the so-called geometrical (multiplicative) convexity. Moreover, it is known that if the function f is differentiable, then it is (a,b)-convex (concave) if and only if $x \mapsto x^{1-a} f'(x)[f(x)]^{b-1}$ is increasing (decreasing). See [3] for more details.

In the next results we can see that the above mentioned (p,q) generalized trigonometric functions are (a,a)-convex/concave. These results refine the earlier results in [6].

Theorem 1. *If* $p, q > 1$ *and* $a \geq 1$, *then* $\operatorname{arcsin}_{p,q}$ *is* (a,a)*-convex on* $(0,1)$, $\operatorname{arctan}_{p,q}$ *is* (a,a)*-concave on* $(0,1)$, *while* $\operatorname{arcsinh}_{p,q}$ *is* (a,a)*-concave on* $(0,\infty)$. *In other words, if* $p,q > 1$ *and* $a \geq 1$, *then we have*

$$\begin{aligned}
\operatorname{arcsin}_{p,q}(M_a(r,s)) &\leq M_a(\operatorname{arcsin}_{p,q}(r), \operatorname{arcsin}_{p,q}(s)), \quad r,s \in (0,1),\\
\operatorname{arctan}_{p,q}(M_a(r,s)) &\geq M_a(\operatorname{arctan}_{p}(r), \operatorname{arctan}_{p}(s)), \quad r,s \in (0,1),\\
\operatorname{arsinh}_{p,q}(M_a(r,s)) &\geq M_a(\operatorname{arsinh}_{p,q}(r), \operatorname{arsinh}_{p,q}(s)), \quad r,s > 0.
\end{aligned}$$

Theorem 2. *If* $p,q > 1$ *and* $a \geq 1$, *then* $\sin_{p,q}$ *is* (a,a)*-concave, and* $\cos_{p,q}$, $\tan_{p,q}$, $\sinh_{p,q}$ *are* (a,a)*-convex on* $(0,1)$. *In other words, if* $p,q > 1$, $a \geq 1$ *and* $r,s \in (0,1)$, *then the next inequalities are valid*

$$\begin{aligned}
\sin_{p,q}(M_a(r,s)) &\geq M_a(\sin_{p,q}(r), \sin_{p,q}(s)),\\
\cos_{p,q}(M_a(r,s)) &\leq M_a(\cos_{p,q}(r), \cos_{p,q}(s)),\\
\tan_{p,q}(M_a(r,s)) &\leq M_a(\tan_{p,q}(r), \tan_{p,q}(s)),\\
\sinh_{p,q}(M_a(r,s)) &\leq M_a(\sinh_{p,q}(r), \operatorname{arsinh}_{p,q}(s)).
\end{aligned}$$

The next theorems improve some of the above results.

Theorem 3. *If* $p,q > 1$, $a \leq 0$ *and* $b \in \mathbb{R}$ *or* $0 < a \leq b$ *and* $b \leq 1$, *then* $\operatorname{arcsin}_{p,q}$ *is* (a,b)*-convex on* $(0,1)$, *and in particular if* $p = q$, *then the function* $\operatorname{arcsin}_p = \operatorname{arcsin}_{p,p}$ *is* (a,b)*-convex on* $(0,1)$. *In other words, if* $p,q > 1$, $a \leq 0$ *and* $b \in \mathbb{R}$ *or* $0 < a \leq b$ *and* $b \leq 1$, *then for all* $r,s \in (0,1)$ *we have*

$$\operatorname{arcsin}_{p,q}(M_a(r,s)) \leq M_b(\operatorname{arcsin}_{p,q}(r), \operatorname{arcsin}_{p,q}(s)).$$

Theorem 4. *If* $p,q > 1$, $a \leq 0 \geq b$ *or* $0 < b \leq a$ *and* $a \leq 1$, *then* $\operatorname{arsinh}_{p,q}$ *is* (a,b)*-concave on* $(0,\infty)$, *and in particular if* $p = q$, *then the function* $\operatorname{arsinh}_p = \operatorname{arsinh}_{p,p}$ *is* (a,b)*-concave on* $(0,\infty)$. *In other words, if* $p,q > 1$, $a \leq 0 \geq b$ *or* $0 < b \leq a$ *and* $a \leq 1$, *then for all* $r,s \in (0,\infty)$ *we have*

$$\operatorname{arsinh}_{p,q}(M_a(r,s)) \geq M_b(\operatorname{arsinh}_{p,q}(r), \operatorname{arsinh}_{p,q}(s)).$$

It is worth to mention that Chu et al. [10] very recently proved implicitly the $(p_1, 0)$-convexity of $\operatorname{arcsin}_{p,q}$ for $p_1 \leq 0$, $p,q > 1$, and the $(p_2, 0)$-concavity of $\operatorname{arcsinh}_{p,q}$ for $p_2 \geq 0$, $p,q > 1$. Our approach is similar but somewhat easier, since we applied in the above cases directly the results from [3]. We also mention that recently Jiang and Qi [13] proved that the $\operatorname{arcsin}_{p,q}$ is $(0,0)$-convex, by using the same idea as we used in this paper. See also [4, 5] for more details.

Now, observe that by using the change of variable $u = (1-t^q)^{1/p}$

$$\operatorname{arccos}_{p,q}(x) = \operatorname{arcsin}_{p,q}(y) = \int_0^y (1-t^p)^{-1/p} dt, \quad \text{where} \quad y = (1-x^p)^{1/q},$$

can be written as

$$\operatorname{arccos}_{p,q}(x) = \frac{p}{q} \int_x^1 f(u) du, \quad \text{where} \quad f(u) = (1-u^p)^{1/q-1} u^{p-2}.$$

Consequently, we have

$$\int_0^x f(u) du = \frac{q}{p} \operatorname{arccos}_{p,q}(0) - \frac{q}{p} \operatorname{arccos}_{p,q}(x) = \frac{q}{p} \operatorname{arcsin}_{p,q}(1) - \frac{q}{p} \operatorname{arccos}_{p,q}(x).$$

The next result is about this integral.

Theorem 5. *The function* $x \mapsto \pi_{p,q}/2 - \operatorname{arccos}_{p,q}(x)$ *is* (a,b)*-convex on* $(0,1)$ *if* $p \in (1,2]$, $q > 1$, $a < 0$ *and* $b \in \mathbb{R}$ *or if* $p,q > 1$, $a \leq 0$ *and* $b \geq 0$. *In other words, if* $p \in (1,2]$, $q > 1$, $a < 0$ *and* $b \in \mathbb{R}$ *or if* $p,q > 1$, $a \leq 0$ *and* $b \geq 0$, *then for all* $r,s \in (0,1)$ *we have*

$$\pi_{p,q}/2 - \operatorname{arccos}_{p,q}(M_a(r,s)) \leq M_b(\pi_{p,q}/2 - \operatorname{arccos}_{p,q}(r), \pi_{p,q}/2 - \operatorname{arccos}_{p,q}(s)).$$

It is worth to mention that if in the above theorem we take $a = b = 0$, then we have that the function $x \mapsto \operatorname{arcsin}_{p,q}(1) - \operatorname{arccos}_{p,q}(x)$ is geometrically convex on $(0,1)$ for $p, q > 1$, and in particular if $p = q$, then the function $x \mapsto \operatorname{arcsin}_{p,p}(1) - \operatorname{arccos}_{p,p}(x)$ is geometrically convex on $(0,1)$ for $p > 1$. In other words, for $p, q > 1$ and $r, s \in (0,1)$ we have

$$\pi_{p,q}/2 - \operatorname{arccos}_{p,q}(\sqrt{rs}) \leq \sqrt{(\pi_{p,q}/2 - \operatorname{arccos}_{p,q}(r))(\pi_{p,q}/2 - \operatorname{arccos}_{p,q}(s))}.$$

2. **Preliminary results**

In this section we present some preliminary results which will be used in the proof of the main theorems. The first known result considers the (p,q)-trigonometric functions whereas the other results consider properties of real functions.

Lemma 1. [12] *For all* $p, q > 1$ *and* $x \in (0, \pi_{p,q}/2)$, *we have*

$$(\sin_{p,q}(x))' = \cos_{p,q}(x),$$

$$(\cos_{p,q}(x))' = -\frac{p}{q}(\cos_{p,q}(x))^{2-p}(\sin_{p,q}(x))^{q-1},$$

$$(\tan_{p,q}(x))' = 1 + \frac{p\,(\sin_{p,q}(x))^q}{q\,(\cos_{p,q}(x))^p}.$$

The following result is known as *the monotone form of l'Hospital's rule.*

Lemma 2. [2, Theorem 1.25] *For* $-\infty < a < b < \infty$, *let* $f, g : [a,b] \to \mathbb{R}$ *be continuous on* $[a,b]$, *and be differentiable on* (a,b). *Let* $g'(x) \neq 0$ *on* (a,b). *If* f'/g' *is increasing (decreasing) on* (a,b), *then so are*

$$x \mapsto \frac{f(x) - f(a)}{g(x) - g(a)} \quad \text{and} \quad x \mapsto \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If f'/g' *is strictly monotone, then the monotonicity in the conclusion is also strict.*

The following two results consider the convexity of the inverse function.

Lemma 3. [15, Theorem 2] *Let* $J \subset \mathbb{R}$ *be an open interval, and let* $f : J \to \mathbb{R}$ *be strictly monotonic function. Let* $f^{-1} : f(J) \to J$ *be the inverse to* f. *If* f *is convex and increasing, then* f^{-1} *is concave.*

Lemma 4. [18, Proposition 2] *Let* $f\colon (a,b) \to (c,d) = f(a,b) \subset \mathbb{R}$ *be a convex function and let* $f^{-1}\colon (c,d) \to \mathbb{R}$ *be its inverse. If* f *is increasing (decreasing), then* f^{-1} *is increasing (decreasing) and concave (convex).*

The following two results consider the (p,q)-convexity of the function.

Lemma 5. [3, Lemma 3] *Let* $a, b \in \mathbb{R}$ *and let* $f\colon [\alpha, \beta] \to (0, \infty)$ *be a differentiable function for* $\alpha, \beta \in (0,\infty)$. *The function* f *is* (a,b)*-convex* ((a,b)*-concave) if and only if* $x \mapsto x^{1-a} f'(x)(f(x))^{b-1}$ *is increasing (decreasing).*

Lemma 6. [3, Theorem 7] *Let* $\alpha, \beta \in (0,\infty)$ *and* $f\colon [\alpha,\beta] \to (0,\infty)$ *be a differentiable function. Denote* $g(x) = \int_\alpha^x f(t)\,dt$ *and* $h(x) = \int_x^\beta f(t)\,dt$. *Then*
(a) If for all $a \in [0,1]$ *the function* f *is* $(a,0)$*-concave, then the function* g *is* (a,b)*-concave for all* $a \in [0,1]$ *and* $b \leq 0$. *If, in addition the function* $x \mapsto x^{1-a} f(x)$ *is increasing for all* $a \in [0,1]$, *then* g *is* (a,b)*-concave for all* $a \in [0,1]$ *and* $b \in (0,1)$. *Moreover, if for all* $a \in \mathbb{R}$ *the function* $x \mapsto x^{1-a} f(x)$ *is increasing, then* g *is* (a,b)*-convex for all* $a \in \mathbb{R}$ *and* $b \geq 1$.
(b) If for all $a \in [0,1]$ *the function* f *is* $(a,0)$*-concave, then the function* g *is* (a,b)*-concave for all* $a \in [0,1]$ *and* $b \leq 0$. *If, in addition the function* $x \mapsto x^{1-a} f(x)$ *is decreasing for all* $a \in [0,1]$, *then* g *is* (a,b)*-concave for all* $a \in [0,1]$ *and* $b \in (0,1)$. *Moreover, if for all* $a \in \mathbb{R}$ *the function* $x \mapsto x^{1-a} f(x)$ *is decreasing, then* g *is* (a,b)*-convex for all* $a \in \mathbb{R}$ *and* $b \geq 1$.

(c) If for all $a \notin (0,1)$ *we have* $\alpha^{1-a}f(\alpha) = 0$ *and the function* f *is* $(a,0)$*-convex, then* g *is* (a,b)*-convex for all* $a \notin (0,1)$ *and* $b \geq 0$*. If, in addition the function* $x \mapsto x^{1-a}f(x)$ *is increasing for all* $a \notin (0,1)$*, then* g *is* (a,b)*-convex for all* $a \notin (0,1)$ *and* $b < 0$*.*
(d) If for all $a \notin (0,1)$ *we have* $\beta^{1-a}f(\beta) = 0$ *and the function* f *is* $(a,0)$*-convex, then* g *is* (a,b)*-convex for all* $a \notin (0,1)$ *and* $b \geq 0$*. If, in addition the function* $x \mapsto x^{1-a}f(x)$ *is decreasing for all* $a \notin (0,1)$*, then* g *is* (a,b)*-convex for all* $a \notin (0,1)$ *and* $b < 0$*.*

Next we introduce a convexity result for the functions $\tan_{p,q}$ and $\operatorname{arctan}_{p,q}$.

Lemma 7. *If* $p,q > 1$*, then the function* $\tan_{p,q}$ *is increasing and convex on* $(0, \pi_{p,q}/2)$*, while the function* $\operatorname{arctan}_{p,q}$ *is increasing and concave on* $(0,1)$*.*

Proof. By Lemma 1 we get

$$[\tan_{p,q}(x)]'' = 1 + \frac{p}{q^2}\left[\frac{pq(\sin_{p,q}(x))^{q-1}(\cos_{p,q}(x))^{1+p} + p^2\cos_{p,q}(x)(\sin_{p,q}(x))^{2q-1}}{(\cos_{p,q}(x))^{2p}}\right],$$

which is positive since both $\sin_{p,q}$ and $\cos_{p,q}$ are positive. Hence $\tan_{p,q}$ is convex. By Lemma 1 we observe that the derivative of $\tan_{p,q}$ is positive and thus $\tan_{p,q}$ is increasing. It follows from Lemma 4 that $\operatorname{arctan}_{p,q}$ is increasing and concave. □

Now, we focus on some monotonicity results.

Lemma 8. *Let* $a \geq 0$*,* $p,q > 1$ *and consider the functions* $f, h : (0,1) \to \mathbb{R}$ *and* $g : (0,\infty) \to \mathbb{R}$*, defined by*

$$f(x) = \left(\frac{\operatorname{arcsin}_{p,q}(x)}{x}\right)^a (\operatorname{arcsin}_{p,q}(x))',$$

$$g(x) = \left(\frac{\operatorname{arsinh}_{p,q}(x)}{x}\right)^a (\operatorname{arsinh}_{p,q}(x))',$$

$$h(x) = \left(\frac{\operatorname{arctan}_{p,q}(x)}{x}\right)^a (\operatorname{arctan}_{p,q}(x))'.$$

Then the function f *is increasing and the functions* g *and* h *are decreasing.*

Proof. It follows from the definition that

$$x \mapsto (\operatorname{arcsin}_{p,q}(x))' = (1 - x^q)^{-1/p},$$

is increasing since

$$(\operatorname{arcsin}_{p,q}(x))'' = \frac{qx^{q-1}(1 - x^q)^{-1-1/p}}{p} > 0.$$

By Lemma 2, the function $x \mapsto (\operatorname{arcsin}_{p,q} x)/x$ is increasing $(0,1)$ and consequently the function $x \mapsto ((\operatorname{arcsin}_{p,q} x)/x)^a$ is increasing too on $(0,1)$. Since the product of increasing functions is increasing, the function f is increasing. Now, the function $\operatorname{arctan}_{p,q}$ is increasing and concave $(0,1)$ by Lemma 7, which implies that $x \mapsto (\operatorname{arctan}_{p,q}(x))'$ is decreasing on $(0,1)$. Using again Lemma 2 it follows that $x \mapsto (\operatorname{arctan}_{p,q}(x))/x$ is decreasing on $(0,1)$, and hence is the function h. Finally, the proof for function g follows similarly as the proof for f, because

$$(\operatorname{arsinh}_{p,q}(x))'' = \frac{-qx^{q-1}(1 + x^q)^{-1-1/p}}{p} < 0.$$ □

Lemma 9. *If* $a \geq 0$ *and* $p,q > 1$*, then the function*

$$x \mapsto \left(\frac{\sin_{p,q}(x)}{x}\right)^a (\sin_{p,q}(x))'$$

is decreasing on $(0,1)$*, while the functions*

$$x \mapsto \left(\frac{\cos_{p,q}(x)}{x}\right)^a (\cos_{p,q}(x))', \; x \mapsto \left(\frac{\tan_{p,q}(x)}{x}\right)^a (\tan_{p,q}(x))'$$

and

$$x \mapsto \left(\frac{\sinh_{p,q}(x)}{x}\right)^a (\sinh_{p,q}(x))'$$

are increasing on $(0,1)$.

Proof. Let $f(x) = \operatorname{arcsin}_{p,q}(x)$, where $x \in (0,1)$. We get

$$f'(x) = \frac{1}{(1-x^q)^{1/p}},$$

which is positive and increasing, hence f is convex. Clearly $\sin_{p,q}$ is increasing, and by Lemma 3 is concave. This implies that $x \mapsto (\sin_{p,q}(x))'$ is decreasing, and $x \mapsto (\sin_{p,q}(x))/x$ is also decreasing by Lemma 2.

Similarly we get that $x \mapsto (\cos_{p,q}(x))'$, $x \mapsto (\tan_{p,q}(x))'$ and $x \mapsto (\sinh_{p,q}(x))'$ are increasing, and the assertion follows from Lemma 2. ☐

3. Proofs of the main results

Proof of the Theorem 1. Let $0 < x < y < 1$, and $u = ((x^a + y^a)/2)^{1/a} > x$. Let us denote $h_1(x) = \operatorname{arcsin}_{p,q}(x)$, $h_2(x) = \operatorname{arctan}_{p,q}(x)$, $h_3(x) = \operatorname{arsinh}_{p,q}(x)$ and for $i \in \{1,2,3\}$ define

$$g_i(x) = h_i(u)^a - \frac{h_i(x)^a + h_i(y)^a}{2}.$$

Differentiating with respect to x, we get $du/dx = (1/2)(x/u)^{a-1}$ and

$$\begin{aligned} g_i'(x) &= \frac{1}{2}\, a\, h_i(x)^{a-1} \frac{d}{dx}(h_i(u)) \left(\frac{x}{u}\right)^{a-1} - \frac{1}{2}\, a\, h_i(x)^{a-1} \frac{d}{dx}(h_i(x)) \\ &= \frac{a}{2} x^{a-1}(f_i(u) - f_i(x)), \end{aligned}$$

where

$$f_i(x) = \left(\frac{h_i(x)}{x}\right)^{a-1} \frac{d}{dx}(h_i(x)).$$

By Lemma 8 g_1' is positive and g_2', g_3' are negative. Hence g_1 is increasing and g_2, g_3 are decreasing. This implies that

$$g_1(x) < g_1(y) = 0, \quad g_2(x) > g_2(y) = 0, \quad g_3(x) > g_3(y) = 0,$$

and the assertion follows. ☐

Proof of the Theorem 2. The proof is similar to the proof of Theorem 1 and follows from Lemma 9. ☐

Proof of Theorem 3. Let us consider the function $f : (0,1) \to (0,\infty)$, $f(t) = (1-t^q)^{-1/p}$. If $a \le 0$, then the function

$$t \mapsto \frac{t^{1-a} f'(t)}{f(t)} = \frac{q}{p}\frac{t^{q-a}}{1-t^q}$$

is increasing on $(0,1)$, that is f is $(a,0)$-convex on $(0,1)$, according to Lemma 5. Since $t^{1-a}(1-t^q)^{-1/p} \to 0$, as $t \to 0$, and $t \to t^{1-a}(1-t^q)^{-1/p}$ is increasing for $a \le 0$ as a product of two increasing and positive functions from Lemma 6, we deduce that $x \mapsto \operatorname{arcsin}_{p,q}(x)$ is (a,b)-convex for $a \le 0$ and $b \ge 0$ or $a \le 0$ and $b < 0$, that is, for $a \le 0$ and $b \in \mathbb{R}$.

According to Theorem 1, the function $x \mapsto \operatorname{arcsin}_{p,q}(x)$ is (a,a)-convex for $a \ge 1$ and $p,q > 1$. Taking into account that the power mean is increasing with respect to its order it follows that $x \mapsto \operatorname{arcsin}_{p,q}(x)$ is (a,b)-convex on $(0,1)$ for $1 \le a \le b$ and $p,q > 1$.

Let $u, v : (0,1) \to (0,\infty)$ be the functions defined by $u(x) = x^{1-a}$ and $v(x) = \varphi(x)(\varphi(x))^{b-1}$ where $\varphi(x) = \operatorname{arcsin}_{p,q}(x)$. Then we have that $u'(x) = (1-a)x^{-a} \ge 0$ if $a \le 1$, $x \in (0,1)$, and

$$v'(x) = (\varphi(x))^{b-2}(\varphi''(x)\varphi(x) + (b-1)(\varphi'(x))^2) \ge 0$$

if $b \geq 1$ and $x \in (0,1)$, since φ is convex. Thus we obtain $x \mapsto x^{1-a}\varphi'(x)(\varphi(x))^{b-1}$ is decreasing on $(0,1)$ if $a \leq 1$ and $b \geq 1$. According to Lemma 5 it follows that $x \mapsto \operatorname{arcsin}_{p,q}(x)$ is (a,b)-convex on $(0,1)$ for $a \leq 1$ and $b \geq 1$.

Finally, we would like to show that the (a,b)-convexity of $x \mapsto \operatorname{arcsin}_{p,q}(x)$ is the case when $b \geq a$ and $b \geq 1$ can be proved by a somewhat different approach. For this, consider the function $\phi : (0,1) \to (0,\infty)$, defined by $\phi(x) = x^{1-a}\varphi'(x)(\varphi(x))^{b-1}$. Then we have

$$\Delta(x) = \frac{x\phi'(x)}{\phi(x)} = 1 - a + \frac{q}{p}\frac{x^q}{1-x^q} + (b-1)\frac{x\varphi'(x)}{\varphi(x)},$$

and

$$\Delta'(x) = \frac{q}{p}\frac{qx^{q-1}}{(1-x^q)^2} + (b-1)\left(\frac{x\varphi'(x)}{\varphi(x)}\right)' \geq 0$$

if $x \in (0,1)$ and $b \geq 1$, since φ is $(0,0)$-convex, as we proved before. Consequently we have $\Delta(x) \geq \Delta(0) = b - a \geq 0$ if $x \in (0,1)$ and $b \geq a$. In other words, if $b \geq a$ and $b \geq 1$, then $x \mapsto \operatorname{arcsin}_{p,q}(x)$ is (a,b)-convex on $(0,1)$. □

Proof of Theorem 4. Observe that for $f : (0,\infty) \to (0,\infty)$, $f(t) = (1+t^q)^{-1/p}$, we have

$$\left(\frac{tf'(t)}{f(t)}\right)' = -\frac{q^2}{p}\frac{t^{q-1}}{(1+t^q)^2} < 0 \quad \text{for all} \quad t > 0 \quad \text{and} \quad p,q > 1.$$

Consequently, f is geometrically concave on $(0,\infty)$ and according to Lemma 6 the function

$$x \mapsto \operatorname{arsinh}_{p,q}(x) = \int_0^x (1+t^q)^{-1/p}dt$$

is also geometrically concave (or (0,0)-concave) on $(0,\infty)$.

Now, consider the functions $u, v, w : (0,\infty) \to (0,\infty)$, defined by

$$u(x) = \frac{x\varphi'(x)}{\varphi(x)}, \quad v(x) = x^{-a}, \quad w(x) = (\varphi(x))^b,$$

where $\varphi(x) = \operatorname{arsinh}_{p,q}(x)$. Observe that for $a \geq 0$ and $b \leq 0$, the function v and w are decreasing, and hence in this case, by using the fact that φ is $(0,0)$-concave, we get that

$$x \mapsto \frac{x\varphi'(x)}{\varphi(x)}x^{-a}(\varphi(x))^b$$

is also decreasing on $(0,\infty)$ as a product of three positive decreasing functions. In other words, φ is (a,b)-concave on $(0,\infty)$ for $a \geq 0$ and $b \leq 0$.

By Theorem 1, the function φ is (a,a)-concave for $a \geq 1$. It follows from the monotonicity of the power mean that φ is (a,b)-concave for $b \leq a$ and $a \geq 1$. □

Proof of Theorem 5. Consider the function $f : (0,1) \to (0,\infty)$, defined by

$$f(t) = (1-t^p)^{1/q-1}t^{p-2}.$$

Then we have

$$\left(\frac{t^{1-a}f'(t)}{f(t)}\right)' = \frac{p}{q}(q-1)\frac{(p-a)t^{p-a-1} + at^{2p-a-1}}{(1-t^p)^2} + a(p-2)t^{-a-1}.$$

If $a = 0$, then we have that f is geometrically convex on $(0,1)$ for $p,q > 1$. This in turn implies that the function

$$x \mapsto \lambda(x) = \frac{q}{p}\operatorname{arcsin}_{p,q}(1) - \frac{q}{p}\operatorname{arccos}_{p,q}(x) = \int_0^x f(t)dt$$

is also geometrically convex on $(0,1)$. Moreover, if $a < 0$, then taking into account that

$$\left(\frac{t^{1-a}f'(t)}{f(t)}\right)' = \frac{p}{q}(q-1)\frac{t^{p-a-1}(p + a(t^p-1))}{(1-t^p)^2} + a(p-2)t^{-a-1},$$

we obtain that for $p \in (1,2]$, $q > 1$ the function f is $(a,0)$-convex. Since $t^{1-a}(1-t^p)^{1/q-1}t^{p-2} \to 0$ as $t \to 0$, according to Lemma 6 the function λ is (a,b)-convex for $a < 0$, $b \geq 0$ and $p \in (1,2]$, $q > 1$. On the other hand, the function

$$t \mapsto t^{1-a}(1-t^p)^{1/q-1}t^{p-2} = t^{p-a-1}(1-t^p)^{1/q-1}$$

is increasing on $(0,1)$ for $p, q > 1$ and $a < 0$. Appealing again to Lemma 6 the function λ is (a,b)-convex for $a < 0$, $b < 0$ and $p \in (1,2]$, $q > 1$.

Finally, if we consider the functions $u, v, w : (0,1) \to (0,\infty)$, defined by

$$u(x) = \frac{x\lambda'(x)}{\lambda(x)}, \quad v(x) = x^{-a}, \quad w(x) = (\lambda(x))^b,$$

then we get the $x \mapsto x^{1-a}\lambda'(x)(\lambda(x))^{b-1}$ is increasing on $(0,1)$ as a product of three positive increasing functions u, v and w when $a \leq 0$ and $b \geq 0$. Here we used that λ is geometrically convex on $(0,1)$. □

REFERENCES

[1] M. Abramowitz and I. Stegun, eds.: *Handbook of mathematical functions with formulas, graphs and mathematical tables.* National Bureau of Standards, 1964 (Russian translation, Nauka 1979).

[2] G.D. Anderson, M.K. Vamanamurthy and M. Vuorinen: *Conformal invariants, inequalities and quasiconformal maps.* J. Wiley, 1997.

[3] Á. Baricz: *Geometrically concave univariate distributions*, J. Math. Anal. Appl. 363(1) (2010) 182–196.

[4] Á. Baricz, B.A. Bhayo and M. Vuorinen: *Turán type inequalities for generalized inverse trigonometric functions*, `arXiv:1305.0938`.

[5] B.A. Bhayo and M. Vuorinen: *Inequalities for eigenfunctions of the p-Laplacian*, `arXiv.1101.3911`.

[6] B.A. Bhayo and M. Vuorinen: *On generalized trigonometric functions with two parameters*, J. Approx. Theory 164 (2012) 1415–1426.

[7] B.A. Bhayo and M. Vuorinen: *Power mean inequality of generalized trigonometric functions*, `arXiv.1209.0873`.

[8] R.J. Biezuner, G. Ercole and E.M. Martins: *Computing the first eigenvalue of the p-Laplacian via the inverse power method*, J. Funct. Anal. 257 (2009) 243–270.

[9] P.J. Bushell and D.E. Edmunds: *Remarks on generalised trigonometric functions*, Rocky Mountain J. Math. 42 (2012) 25–57.

[10] Y.-M. Chu, Y.-P. Jiang and M.-K. Wang: *Inequalities for generalized trigonometric and hyperbolic sine functions*, `arXiv.1212.4681`.

[11] P. Drábek and R. Manásevich: *On the closed solution to some p−Laplacian nonhomogeneous eigenvalue problems.* Differential Integral Equations 12 (1999) 773–788.

[12] D.E. Edmunds, P. Gurka and J. Lang: *Properties of generalized trigonometric functions*, J. Approx. Theory 164 (2012) 47–56.

[13] W.-D. Jiang and F. Qi: *Geometric convexity of the generalized sine and the generalized hyperbolic sine*, `arXiv.1301.3264`.

[14] R. Klén, V. Manojlović, S. Simić and M. Vuorinen: *Bernoulli inequality and hypergeometric functions*, Proc. Amer. Math. Soc. (in press).

[15] M. Kuczma: *An introduction to the theory of functional equations and inequalities.* Cauchy's equation and Jensen's inequality, With a Polish summary. Prace Naukowe Uniwersytetu Ślaskiego w Katowicach [Scientific Publications of the University of Silesia], 489. Uniwersytet Ślaski, Katowice; Państwowe Wydawnictwo Naukowe (PWN), Warsaw, 1985.

[16] J. Lang and D.E. Edmunds: *Eigenvalues, Embeddings and Generalised Trigonometric Functions*, Lecture Notes in Mathematics, vol. 2016, Springer-Verlag, 2011.

[17] P. Lindqvist: *Some remarkable sine and cosine functions*, Ricerche di Matematica 44 (1995) 269–290.

[18] M. Mršević: *Convexity of the inverse function*, The teaching of mathematics 11 (2008) 21–24.

[19] S. Takeuchi: *Generalized Jacobian elliptic functions and their application to bifurcation problems associated with p-Laplacian*, J. Math. Anal. Appl. 385 (2012) 24–35.

Department of Economics, Babeş-Bolyai University, 400591 Cluj-Napoca, Romania
E-mail address: `bariczocsi@yahoo.com`

Department of Mathematical Information Technology, University of Jyväskylä, 40014 Jyväskylä, Finland
E-mail address: `bhayo.barkat@gmail.com`

Department of Mathematics and Statistics, University of Turku, 20014 Turku, Finland
E-mail address: `riku.klen@utu.fi`